

J. Benjamin McIntire, M.D. 537 Stanton-Christiana Rd., Newark, DE 19713

537 Stanton-Christiana Rd., Newark, DE 19713 P: 302-998-8007 F: 302-225-3775

www.premiercosmeticsurgeryde.com

to take

Preop Checklist

At the time of surgical scheduling:

	If your procedure is under oral sedation, IV sedation, or general anesthesia, please arrange for someone you home from the hospital or surgery center. You will not be allowed to drive or leave alone.						
6	wee	ks	before surgery:				
	Dis	Discontinue all nicotine products					
2 .	wee	ks	before surgery:				
	Dis	Discontinue all medications/supplements that may thin your blood:					
	- - - -	Ibu Naj Any Vita Her	pirin profen (Advil, Motrin) proxen (Aleve) y other non-steroidal anti-inflammatory medications amin E rbal supplements y other medications containing any of these compounds				
	Discontinue GLP-1 inhibitors (Ozempic class)						
	If undergoing septorhinoplasty, discontinue steroid nasal sprays (Flonase, Nasacort, etc)						
	Pre	par	e for recovery and wound care (refer to your specific procedure):				
	- The following items are beneficial for all types of surgery :						
			Tylenol, regular strength – to be used instead of (not in addition to) narcotic pain medication				
			Probiotics – to help reduce GI distress from oral antibiotics				
			Ice packs – to help reduce bruising/swelling in the early period				
			Wedge pillow – to keep head elevated while sleeping				
			Throat spray/lozenges – to help with sore throat after sedation/general anesthesia				
	-	If u	indergoing septorhinoplasty:				
			Q-tips				
			Hydrogen peroxide				
			Nasal saline spray				

			Eyeglass suspenders			
	-	If undergoing eyelid surgery:				
			Q-tips			
			Distilled water			
			Aquaphor			
- If undergoing any other skin surgery (facelift, submentoplasty, otoplasty, earlobe repa skin cancer repair, cyst/lipoma excision, etc):			indergoing any other skin surgery (facelift, submentoplasty, otoplasty, earlobe repair, scar revision, n cancer repair, cyst/lipoma excision, etc):			
			Q-tips			
			Hydrogen peroxide			
			Vaseline			
			4x4 gauze			
			Telfa "nonstick" pads			
			3M micropore tape			
	Pre	e for bruising (optional):				
		Bro	omelain – 500mg twice per day, starting 1 week before surgery and continuing 2 weeks after surgery			
		Arr	nica – 4 pills 4 times per day, starting 4 days before surgery and continuing 4 days after surgery			
1 day before surgery:						
	Ph	Phone call from the hospital or surgery center to give you your arrival time for surgery				
	Sh	Shower at night, no cologne or perfume				
	Ifι	If undergoing IV sedation or general anesthesia, do NOT eat or drink anything, including water, after midnigh				
Sı	ırg	ery	day:			
	We	Wear loose clothing, no makeup, no piercings				
	Bri	Bring ID, insurance card, and containers for glasses/contacts/dentures/hearing aids				